

ENVIRONMENTAL TOXICITY OF THE EXPLOSIVES RDX AND TNT IN SOIL TO THE SOIL INVERTEBRATE *FOLSOMIA CANDIDA*

Carlton T. Phillips, Ronald T. Checkai, Roman G. Kuperman, Michael Simini,
Jan E. Kolakowski, and Carl W. Kurnas
U.S. Army Edgewood Chemical Biological Center (ECBC)
Aberdeen Proving Ground, MD 21010-5424 USA

ABSTRACT

We investigated the toxicity of hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX) and 2,4,6-trinitrotoluene (TNT) to the soil invertebrate *Folsomia candida* (Collembola) in five natural soils: Sassafras sandy loam (SSL), Teller sandy loam (TSL), Richfield clay loam (RCL), Kirkland clay loam (KCL), and Webster clay loam (WCL). According to the Ecological Soil Screening Level (Eco-SSL) criteria (USEPA 2003), relative bioavailability scores for organic chemicals in these soils were considered “high” for SSL and TSL; “medium” for RCL and KCL; and relatively “low” for WCL soil. We investigated whether soil type affects the toxicity of RDX or TNT in soil to Collembola by adapting a standardized *Folsomia* reproduction test (ISO 11267:1998). Adult survival and juvenile production by *F. candida* were assessed using replicated multiple treatment concentrations and appropriate controls. Juvenile production data were analyzed using nonlinear regression models. Initial EC₂₀ and EC₅₀ values [mg kg⁻¹] for juvenile production in freshly amended soils were, respectively for RDX: 7 and 30 (TSL), 16 and 55 (RCL), 28 and 90 (SSL), 31 and 110 (KCL), 3 and 50 (WCL); and for TNT: 7 and 17 (TSL), 4 and 23 (RCL), 17 and 25 (SSL), 21 and 37 (KCL), 174 and 259 (WCL). These provisional results indicate that the soil with highest organic matter and clay contents (WCL) sustained the least TNT toxicity, while soil with lowest organic matter and clay contents (TSL), sustained the greatest TNT toxicity to *F. candida* reproduction based on EC₅₀ values. Correlation analysis showed relatively weak relationships overall among the soil properties and toxicity endpoints for RDX.

1. INTRODUCTION

Objective Force Soldiers must be highly trained in all tasks across the spectrum of military operations. These Soldiers will need demanding, highly realistic training to obtain this capability. This increased training will lead to an increased release into the environment of explosives at training sites. Soil contamination was identified at more than 21,000

sites among Department of Defense (DoD) installations (Bridges and Whaley, 1997). By 2001, the number of known waste sites on current and former DoD installations in the United States exceeded 28,000. Many of these sites associated with military operations that involve munition manufacturing, disposal, testing, and training, contain elevated levels of explosives and related materials in soil.

We adapted the standardized Collembola Reproduction Test to measure the toxicity of RDX and TNT to this ecologically relevant test species. This chronic assay includes at least one reproductive component among the measurement endpoints.

Soils with a relatively wide range of physical/chemical characteristics were tested, including: Sassafras sandy loam (SSL), Teller sandy loam (TSL), Richfield clay loam (RCL), Kirkland clay loam (KCL), and Webster clay loam (WCL).

2. RESULTS AND DISCUSSION

The effect of RDX or TNT on adult survival of *F. candida* varied among freshly amended soil types tested. RDX toxicity measured as adult *F. candida* mortality was highest in TSL and lowest in SSL. The greatest TNT toxicity to adult *F. candida* was found in the soil with lowest organic matter and clay contents (TSL), and was lowest in the soil with highest organic matter and clay contents (WCL).

The EC₂₀ and EC₅₀ values (mg kg⁻¹) for juvenile production in freshly amended soils were, respectively for RDX: 7 and 30 (TSL), 16 and 55 (RCL), 28 and 90 (SSL), 31 and 110 (KCL), 3 and 50 (WCL); and for TNT: 7 and 17 (TSL), 4 and 23 (RCL), 17 and 25 (SSL), 21 and 37 (KCL), 174 and 260 (WCL).

Preliminary results based on EC₅₀ values indicate that the soil with highest organic matter and clay contents (WCL) sustained the least RDX or TNT toxicity to *F. candida* reproduction, while soil with lowest organic matter and clay contents (TSL)

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 00 DEC 2004		2. REPORT TYPE N/A		3. DATES COVERED -	
4. TITLE AND SUBTITLE Environmental Toxicity Of The Explosives Rdx And Tnt In Soil To The Soil Invertebrate Folsomia Candida				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. Army Edgewood Chemical Biological Center (ECBC) Aberdeen Proving Ground, MD 21010-5424 USA				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited					
13. SUPPLEMENTARY NOTES See also ADM001736, Proceedings for the Army Science Conference (24th) Held on 29 November - 2 December 2005 in Orlando, Florida. , The original document contains color images.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 2	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

sustained the greatest RDX or TNT toxicity to *F. candida* reproduction. However, the soils having intermediate levels of organic matter and clay contents were not as consistent in their modification of the resulting RDX or TNT toxicities to *F. candida*.

ACKNOWLEDGEMENT

This work was supported by the Strategic Environmental Research and Development Program.

REFERENCES

- Bridges, T.S. and Whaley, J.E. (1997). *Summary of a workshop on ecological risk assessment and military-related compounds: Current research needs*. US Army Corps of Engineers, Waterways Experiment Station, Miscellaneous Paper IRRP-97-1. March 1997, 72 pp.
- ISO (International Standardization Organization) (1998b). *Soil Quality – Inhibition of Reproduction of Collembola (Folsomia candida) by Soil Pollutants*. ISO 11267:1998(E).
- USEPA. (2003). *Ecological Soil Screening Level Guidance*. U.S. Environmental Protection Agency. Office of Emergency and Remedial Response. Washington, DC.

CONCLUSION

The overall results indicate that soil physical and chemical properties may alter the toxicity of RDX and TNT. This is shown by different dose responses by *F. candida* exposed to the same concentrations of either RDX or TNT in soils having different characteristics. However, based on the relative differences in the modification of RDX or TNT toxicities to *F. candida* in soils having intermediate levels of organic matter and clay contents, ecological assessments of risk should not be based solely upon any single bioassay.